


# The Carbon-Land Model Intercomparison Project (C-LAMP)

Forrest M. Hoffman<sup>1</sup>, James T. Randerson<sup>2</sup>,  
Peter E. Thornton<sup>3,1</sup>, Natalie M. Mahowald<sup>3,4</sup>, Keith Lindsay<sup>3</sup>,  
Yen-Huei Lee<sup>3</sup>, Cynthia D. Nevison<sup>5,3</sup>, Scott C. Doney<sup>6</sup>,  
Gordon B. Bonan<sup>3</sup>, Reto Stöckli<sup>7,8</sup>, Curtis C. Covey<sup>9</sup>,  
Steven W. Running<sup>10</sup>, and Inez Y. Fung<sup>11</sup>

<sup>1</sup>Oak Ridge National Laboratory, <sup>2</sup>University of California-Irvine, <sup>3</sup>National Center for Atmospheric Research, <sup>4</sup>Cornell University, <sup>5</sup>University of Colorado-Boulder, <sup>6</sup>Woods Hole Oceanographic Institute, <sup>7</sup>Colorado State University, <sup>8</sup>MeteoSwiss, <sup>9</sup>Lawrence Livermore National Laboratory/PCMDI, <sup>10</sup>University of Montana, and <sup>11</sup>University of California-Berkeley

ANU Env. Biology Seminar • 21 August 2009 • Canberra, AU

- The **Carbon-Land Model Intercomparison Project (C-LAMP)** began as a **CCSM Biogeochemistry Working Group** project to assess model capabilities in the coupled climate system and to explore processes important for inclusion in the CCSM4 Earth System Model for use in the IPCC Fifth Assessment Report (AR5).
- Unlike traditional MIPs, C-LAMP was designed to confront models with best-available observational datasets, develop metrics for evaluation of biosphere models, and build a general-purpose BGC diagnostics package for model evaluation.




- C-LAMP is a Biogeochemistry Subproject of the **Computational Climate Science End Station** (Warren Washington, PI), a U.S. Dept. of Energy INCITE Project.
- Models were initially run on the Cray X1E vector supercomputer in ORNL's **National Center for Computational Sciences (NCCS)**.  
**Cray X1E (phoenix)**



1024 processors (MSPs), 2048 GB memory, and 18.08 TFlop/s peak  
**DECOMMISSIONED September 30, 2008**

# Present Jaguar: 250 TFlop/s





# New Jaguar: Second Fastest in the World at 1.059 PFlop/s



**World's Most Powerful Computer.  
For Science!**

"The Jaguar system at ORNL provides immense computing power in a balanced, stable system that is allowing scientists and engineers to tackle some of the world's most challenging problems."  
—2008, Kelvin Droegemeier, Meteorology Professor, University of Oklahoma.

# C-LAMP Protocol Overview

- **Experiment 1:** Models forced with an improved NCEP/NCAR reanalysis climate data set (Qian, *et al.* 2006) to examine the influence of climate variability, prescribed atmospheric CO<sub>2</sub>, and land cover change on terrestrial carbon fluxes during the 20th century (specifically 1948–2004).
- **Experiment 2:** Models coupled with an active atmosphere (CAM3), prescribed atmospheric CO<sub>2</sub>, prescribed sea surface temperatures and ocean carbon fluxes to examine the effect of a coupled biosphere-atmosphere for carbon fluxes and climate during the 20th century.
- CCSM3.1 partially coupled (“I” & “F” configurations) run at T42 resolution ( $\sim 2.8^\circ \times 2.8^\circ$ ), spectral Eulerian dycore,  $1^\circ \times 0.27^\circ$ – $0.53^\circ$  ocean & sea ice data models (T42gx1v3).
- Experimental protocol, output fields, and metrics are available at <http://www.climatemodeling.org/c-lamp/>

# C-LAMP, C<sup>4</sup>MIP, and iLEAPS

- C-LAMP Experiment 2 is patterned after C<sup>4</sup>MIP (Coupled Climate-Carbon Cycle Model Intercomparison Project, <http://www.c4mip.cnrs-gif.fr/>) Phase 1.
- At the October 2006 C<sup>4</sup>MIP Workshop at the UK Met Office in Exeter, there was strong interest in Experiment 1 and validation experiments using Fluxnet observations.
- At the Marie Curie/iLEAPS Workshop in Hyères, a number of modeling groups expressed interest in consistent model validation and model-data comparisons for their coupled biosphere models, but best-available observations from ground and satellite measurements are difficult to manipulate.
- C-LAMP is sharing forcing and observational datasets, and model results are available through the Earth System Grid (ESG), just like for CMIP3 (the IPCC AR4 model results).

*Offline Forcing with NCEP/NCAR Reanalysis*

Exp.	Description	Time Period
1.1	Spin Up	~4,000 y
1.2	Control	1798–2004
1.3	Varying climate	1948–2004
1.4	Varying climate, CO <sub>2</sub> , and N deposition	1798–2004
1.5	Varying climate, CO <sub>2</sub> , N deposition and land use	1798–2004
1.6	Free Air CO <sub>2</sub> Enrichment (FACE) Control	1997–2100
1.7	Free Air CO <sub>2</sub> Enrichment (FACE) Transient	1997–2100

*Coupled Land-Atmosphere Forcing with Hadley SSTs*

Exp.	Description	Time Period
2.1	Spin Up	~2,600 y
2.2	Control	1800–2004
2.3	Varying climate	1800–2004
2.4	Varying climate, CO <sub>2</sub> , and N deposition	1800–2004
2.5	Varying climate, CO <sub>2</sub> , N deposition and land use	1800–2004
2.6	Varying climate, CO <sub>2</sub> , N deposition, seasonal FFE	1800–2004

All but the land use experiments were run with CCSM3.1  
using CLM3-CASA' and CLM3-CN biogeochemistry models  
yielding >16,000 y and ~50 TB

**C-LAMP Common Model Output**

While all models participating in the Carbon Land Model intercomparison Project (C-LAMP) will output their own "native" fields, a common set of fields is needed to facilitate head-to-head comparison of the models to each other and to available observational datasets. Model results transmitted to the [Earth System Grid](#) for redistribution to the community will use common field names, netCDF long names, [CF Standard Names](#) and units. Contained below is a table of the common output fields required for the C-LAMP and consistent with the metadata conventions used for [CMIP3](#), formerly called the IPCC 4<sup>th</sup> Assessment Model Output database. Corrections and suggestions are solicited on this information. Software is available for rewriting model output into netCDF files following the [Climate and Forecast \(CF\) Metadata Convention](#).

Version 2.1 - Aug 30, 2008

Atmospheric forcing				
Variable Name	Long Name and CF Standard Name	Units	Comment	Statistics
husf	Specific humidity at atmospheric forcing height specific_humidity†	kg kg <sup>-1</sup>		MHM, MHS, MM
prra	Rainfall precipitation flux rainfall_flux†	kg m <sup>-2</sup> s <sup>-1</sup>	Rainfall includes all liquid types (rain, large-scale, convective, etc.)	MHM, MHS, MM
prsn†	Snowfall precipitation flux snowfall_flux†	kg m <sup>-2</sup> s <sup>-1</sup>	Snowfall includes all frozen types (snow, hail, ice, etc.)	MHM, MHS, MM
Biogeochemistry				
Variable Name	Long Name and CF Standard Name	Units	Comment	Statistics
agbc*	Above-ground biomass carbon above_ground_biomass_carbon_content	kg m <sup>-2</sup>	Total carbon content in above-ground live and dead carbon pool(s)	MM
aglbc*	Above-ground live biomass carbon above_ground_live_biomass_carbon_content	kg m <sup>-2</sup>	Total carbon content in above-ground live carbon pool(s)	MM
agnpp	Above-ground net primary production above_ground_net_primary_productivity_of_carbon	kg m <sup>-2</sup> s <sup>-1</sup>	Component of net primary production attributable to above-ground live biomass	MM
ar	Autotrophic respiration autotrophic_respiration_of_carbon alias(es): plant_respiration_carbon_flux	kg m <sup>-2</sup> s <sup>-1</sup>	Sum of maintenance respiration and growth respiration of vegetation	MHM, MHS, MM
bco	Biogenic carbon monoxide flux biogenic_carbon_monoxide_flux	kg m <sup>-2</sup> s <sup>-1</sup>	Total biogenic carbon monoxide flux out of biosphere	MM

Done

# C-LAMP Performance Metrics and Diagnostics

- An evolving draft document on metrics for model evaluation is available at <http://www.climatemodeling.org/c-lamp/>
- Each model is scored with respect to its performance on various output fields compared with best-available observational datasets.
- Examples include:
  - net primary production (NPP) from EMDI and MODIS
  - leaf area index (LAI) using MODIS spatial distribution and phase
  - CO<sub>2</sub> seasonal cycle (NOAA/Globalview flask sites, after running fluxes through an atmospheric transport model for Experiment 1)
  - regional carbon stocks (Saatchi *et al.*, 2006; Batjes, 2006)
  - carbon and energy fluxes (Fluxnet sites)
  - transient dynamics ( $\beta$  factor, etc.)
- More diagnostic or metric ideas? Please contribute them!

Score Sheet for CLAMP - Mozilla Firefox

http://www.climate modeling.org/c-lamp/results/diagnostics/CN\_vs\_CN/

### C-LAMP Score Sheet for Biogeochemical Model Evaluation


Metric	Metric components	Observations & comparison protocol	Model CASA'	Model CN	Score (points)		
					Possible	CASA'	CN
LAI	MODIS Phase	<a href="#">global map</a>	<a href="#">global map model vs obs</a>	<a href="#">global map model vs obs</a>	6.00	5.11	4.24
	MODIS Maximum	<a href="#">global map</a>	<a href="#">global map model vs obs</a>	<a href="#">global map model vs obs</a>	5.00	4.60	4.26
	MODIS Mean	<a href="#">land class obs</a> <a href="#">land class model</a> <a href="#">global map</a>	<a href="#">model vs obs table</a> <a href="#">global map model vs obs</a>	<a href="#">model vs obs table</a> <a href="#">global map model vs obs</a>	4.00	3.75	3.53
NPP	EMDI NPP observations	<a href="#">Class A table</a>	<a href="#">table scatter plot</a>	<a href="#">table scatter plot</a>	1.00	0.68	0.73
		<a href="#">Class B table</a>	<a href="#">table scatter plot</a>	<a href="#">table scatter plot</a>	1.00	0.83	0.82
	EMDI NPP normalized by PPT	<a href="#">Class A histogram</a>	<a href="#">Class A histogram</a>	<a href="#">Class A histogram</a>	2.00	1.50	1.74
		<a href="#">Class B histogram</a>	<a href="#">Class B histogram</a>	<a href="#">Class B histogram</a>	2.00	1.51	1.65
	Correlation with MODIS	<a href="#">global map</a>	<a href="#">model map model vs obs</a>	<a href="#">model map model vs obs</a>	2.00	1.64	1.44
Correlation with MODIS-zonal mean	<a href="#">zonal mean obs</a>	<a href="#">zonal mean model vs obs plot</a>	<a href="#">zonal mean model vs obs plot</a>	2.00	1.88	1.84	
CO <sub>2</sub> Seasonal Cycle — Comparison with Globalview phase and amplitude	60°N–90°N	—	—	—	6.00	4.11	2.77
	30°N–60°N	—	—	—	6.00	4.23	3.23
	0°N–30°N	—	—	—	3.00	2.07	1.71
Energy and C Fluxes from Fluxnet	NEE	—	—	—	—	—	—
	Net radiation	—	—	—	—	—	—
	Latent heat	<a href="#">line plot</a>	<a href="#">model vs obs</a>	<a href="#">model vs obs</a>	—	—	—
Energy and C Fluxes from Ameriflux	Sensible heat	—	—	—	—	—	—
	NEE	—	—	—	6.00	2.46	2.13
	Shortwave Incoming	—	—	—	—	—	—
	Latent heat	<a href="#">line plot</a>	<a href="#">model vs obs timeseries plot</a>	<a href="#">model vs obs timeseries plot</a>	9.00	6.38	6.39
	Sensible heat	—	—	—	9.00	4.90	4.64
	GPP	—	—	—	6.00	3.39	3.46
Aboveground live biomass in South America	ER	—	—	—	—	—	—
	Amazon	<a href="#">obs amazon</a>	<a href="#">model amazon</a>	<a href="#">amazon map</a>	10.00	5.20	4.00

Score Sheet for CLAMP - Mozilla Firefox								
File Edit View History Bookmarks Tools Help								
http://www.climate modeling.org/c-lamp/results/diagnostics/CN_vs_C/ Google								
CO <sub>2</sub> Seasonal Cycle — Comparison with Globalview phase and amplitude	60°N–90°N	—	—	—	6.00	4.11	2.77	
	30°N–60°N	—	—	—	6.00	4.23	3.23	
	0°N–30°N	—	—	—	3.00	2.07	1.71	
Energy and C Fluxes from Fluxnet	NEE	—	—	—	—	—	—	
	Net radiation	—	—	—	—	—	—	
	Latent heat	<a href="#">line plot</a>	<a href="#">model vs obs</a>	<a href="#">model vs obs</a>	—	—	—	
	Sensible heat	—	—	—	—	—	—	
Energy and C Fluxes from Ameriflux	NEE	—	—	—	6.00	2.46	2.13	
	Shortwave Incoming	—	—	—	—	—	—	
	Latent heat	<a href="#">line plot</a>	<a href="#">model vs obs timeseries plot</a>	<a href="#">model vs obs timeseries plot</a>	9.00	6.38	6.39	
	Sensible heat	—	—	—	9.00	4.90	4.64	
	GPP	—	—	—	6.00	3.39	3.46	
	ER	—	—	—	—	—	—	
Carbon Stocks and Transient Dynamics	Aboveground live biomass in South America	<a href="#">obs amazon</a>	<a href="#">model amazon model vs obs</a>	<a href="#">amazon map model vs obs</a>	10.00	5.28	4.99	
	Aboveground live biomass within Amazon Basin (sum within Legal Amazon)	<a href="#">mask obs masked 68.90 (Pg C)</a>	<a href="#">model masked model vs obs 198.87 (Pg C)</a>	<a href="#">model masked model vs obs 160.61 (Pg C)</a>	—	—	—	
	NPP Stimulation from elevated CO <sub>2</sub>	—	<a href="#">FACE Site table biome table</a>	<a href="#">FACE Site table biome table</a>	10.00	7.87	4.11	
	Interannual variability of global carbon fluxes - comparison with TRANSCOM	—	—	—	5.00	3.55	3.00	
	Turnover times and pool sizes	—	<a href="#">Leaf Wood Fine Root Litter Coarse Woody Debris Soil</a>	<a href="#">Leaf Wood Fine Root Litter Coarse Woody Debris Soil</a>	—	—	—	
	Carbon Sinks (1990–2004)	—	<a href="#">biome mean biome total</a>	<a href="#">biome mean biome total</a>	—	—	—	
	Fire Variability (1997–2004)	—	—	<a href="#">global spatial comparison temporal dynamics</a>	5.00	—	1.70	
	<b>Total Score</b>				<b>100.00</b>	<b>65.74</b>	<b>58.38</b>	

Done




- Comparisons with field observations include net primary production (NPP) from the Ecosystem Model-Data Intercomparison (EMDI).
- Measurements were performed in different ways, at different times, and by different groups for a limited number of field sites.
- Shown here are comparisons of NPP with EMDI Class A observations (Figures a and b) and Class B observations (Figures c and d).




*Data provided by NASA Distributed Active Archive Center (DAAC) at ORNL*


- Comparisons with satellite “modeled observations” must be made carefully because of high uncertainty.
- This comparison with MODIS leaf area index (LAI) focuses on the month of maximum LAI (phase), a measurement with less uncertainty than the “observed” LAI values.
- C-LAMP accounts for this uncertainty by weighting scores accordingly.
- CLM-CASA' scored 5.1/6.0 while CLM-CN scored 4.2/6.0 for this metric.




- MODIS net primary production (NPP) “observations” have higher uncertainty.
- Comparison with MODIS NPP focuses on correlation of spatial patterns.
- CLM-CASA' scored 1.6/2.0 while CLM-CN scored 1.4/2.0.





- Estimates of carbon stocks are very difficult to obtain.
- This comparison with estimates of aboveground live biomass in the Amazon by Saatchi *et al.* (2006) shows that both models are too high by about a factor of 2.
- Using a score based on normalized cell-by-cell differences, CLM-CASA' scored 5.3/10.0 while CLM-CN scored 5.0/10.0.




- Comparisons with AmeriFlux eddy correlation CO<sub>2</sub> flux tower sites include net ecosystem exchange (NEE), gross primary production (GPP), respiration, shortwave incoming radiation, and latent and sensible heat.
- Shown here is a comparison of CLM-CASA' results with the Morgan Monroe L4 time series data.
- All AmeriFlux data are stored and distributed by ORNL's Carbon Dioxide Information Analysis Center (CDIAC).



- Additional field measurement comparisons include the Free Air CO<sub>2</sub> Enrichment (FACE) results, including the ORNL site.
- The Norby *et al.* (2005) synthesis of four FACE site observations suggested “response of forest NPP to elevated [CO<sub>2</sub>] is highly conserved across a broad range of productivity, with a stimulation at the median of  $23 \pm 2\%$ .”
- A C-LAMP experiment was added to test this result by increasing [CO<sub>2</sub>] to 550 ppmv in 1997.





Site Name	Lon (°E)	Lat (°N)	Observations		CASA'			CN		
			NPP↑	$\beta_L$	NPP↑	$\beta_L$	Score	NPP↑	$\beta_L$	Score
Duke	-79.08	35.97	28.0%	0.69	16.4%	0.41	0.26	6.2%	0.15	0.65
Aspen	-89.62	45.67	35.2%	0.87	15.6%	0.39	0.39	12.4%	0.31	0.48
ORNL	-84.33	35.90	23.9%	0.59	17.3%	0.43	0.16	5.2%	0.13	0.64
POP-Euro	11.80	42.37	21.8%	0.54	20.0%	0.49	0.04	5.7%	0.14	0.59
4 site mean			27.2%	0.67	17.3%	0.43		7.4%	0.18	
Total M Score							0.79	0.41		

**But!** Norby is now reporting reduced NPP enhancement at the ORNL FACE site due probably to N limitation!


# C-LAMP Score Sheet for CLM3-CASA' and CLM3-CN

Metric	Metric components	Uncertainty of obs.	Scaling mismatch	Total score	Sub-score	CASA'	CN
LAI	Matching MODIS observations			15.0		13.5	12.0
	• Phase (assessed using the month of maximum LAI)	Low	Low		6.0	5.1	4.2
	• Maximum (derived separately for major biome classes)	Moderate	Low		5.0	4.6	4.3
	• Mean (derived separately for major biome classes)	Moderate	Low		4.0	3.8	3.5
NPP	Comparisons with field observations and satellite products			10.0		8.0	8.2
	• Matching EMDI Net Primary Production observations	High	High		2.0	1.5	1.6
	• EMDI comparison, normalized by precipitation	Moderate	Moderate		4.0	3.0	3.4
	• Correlation with MODIS ( $r^2$ )	High	Low		2.0	1.6	1.4
	• Latitudinal profile comparison with MODIS ( $r^2$ )	High	Low		2.0	1.9	1.8
CO <sub>2</sub> annual cycle	Matching phase and amplitude at Globalview flash sites			15.0		10.4	7.7
	• 60°–90°N	Low	Low		6.0	4.1	2.8
	• 30°–60°N	Low	Low		6.0	4.2	3.2
	• 0°–30°N	Moderate	Low		3.0	2.1	1.7
Energy & CO <sub>2</sub> fluxes	Matching eddy covariance monthly mean observations			30.0		17.2	16.6
	• Net ecosystem exchange	Low	High		6.0	2.5	2.1
	• Gross primary production	Moderate	Moderate		6.0	3.4	3.5
	• Latent heat	Low	Moderate		9.0	6.4	6.4
	• Sensible heat	Low	Moderate		9.0	4.9	4.6
Transient dynamics	Evaluating model processes that regulate carbon exchange on decadal to century timescales			30.0		16.8	13.8
	• Aboveground live biomass within the Amazon Basin	Moderate	Moderate		10.0	5.3	5.0
	• Sensitivity of NPP to elevated levels of CO <sub>2</sub> : comparison to temperate forest FACE sites	Low	Moderate		10.0	7.9	4.1
	• Interannual variability of global carbon fluxes: comparison with TRANSCOM	High	Low		5.0	3.6	3.0
	• Regional and global fire emissions: comparison to GFEDv2	High	Low		5.0	0.0	1.7
					<b>Total: 100.0</b>		<b>65.9</b>



## Collaboration with SciDAC Visualization and Analytics Center for Enabling Technologies (VACET)


- C-LAMP and other model results are being used by members of VACET at the National Center for Computational Sciences (NCCS) to explore high performance visualization techniques.




# Collaboration with SciDAC Visualization and Analytics Center for Enabling Technologies (VACET)

- C-LAMP model results and MODIS satellite data are being used by Jian Huang's group at the University of Tennessee, Knoxville (UTK) applying novel statistical methods to the analysis of very large climate data sets.

CLM3-CASA' C-LAMP Control



CLM3-CASA' C-LAMP Transient



The slope of temporal change ( $\lambda$ ) in exposed one-sided leaf area index (ELAI) relative to the April–May change. Red areas “green up” sooner in the year while blue areas “green up” later in the year over the 1850–2000 period.

# Earth System Grid (ESG) Node at ORNL for C-LAMP

C-LAMP Model Data - Mozilla Firefox

File Edit View Go Bookmarks Tools Help

https://esg2.ornl.gov:8443/

**C-LAMP Model Data**

Home Data About ESG Login

Contact ESG

Scientific Discovery through Advanced Computing

**Welcome**

Welcome to the CCES C-LAMP data portal. If you are new to this site, please review the help pages:

[Registration](#)  
[Searching](#)  
[Browsing and Downloading Data](#)  
[Downloading from FTP](#)

**Data Search**

Search Dataset metadata for:

Example: mri, cccma  
[Advanced Search](#)

**Browse Dataset Catalogs**

CCSM Carbon LAnd Model Intercomparison Project (C-LAMP)

CCES C-LAMP Portal Collaborators

[Home](#) | [Data](#) | [About ESG](#) | [Login](#)

Login Status: Not logged in.

© 2004, UCAR. All rights reserved.  
 Portions © 2004, The Regents of the University of California. All rights reserved.  
[Privacy & Security Notices](#)

Done esg2.ornl.gov:8443

# Biases and Weaknesses Exposed by the C-LAMP Analysis

- Both models had a low LAI bias in boreal and arctic regions. This bias was partially eliminated by a new hydrology model capturing freeze-thaw dynamics.

# Biases and Weaknesses Exposed by the C-LAMP Analysis

- Both models had a low LAI bias in boreal and arctic regions. This bias was partially eliminated by a new hydrology model capturing freeze-thaw dynamics.
- Both models had a 1–3 month delay in the timing of maximum LAI. This bias was reduced in CLM3-CN where it was most significant.

# Biases and Weaknesses Exposed by the C-LAMP Analysis

- Both models had a low LAI bias in boreal and arctic regions. This bias was partially eliminated by a new hydrology model capturing freeze-thaw dynamics.
- Both models had a 1–3 month delay in the timing of maximum LAI. This bias was reduced in CLM3-CN where it was most significant.
- Both models overestimate woody biomass in the Amazon Basin. Carbon comparisons with Malhi *et al.* (in press) suggest too much allocation to wood. Allocation in CLM3-CN was adjusted to reduce this bias.

# Biases and Weaknesses Exposed by the C-LAMP Analysis

- Both models had a low LAI bias in boreal and arctic regions. This bias was partially eliminated by a new hydrology model capturing freeze-thaw dynamics.
- Both models had a 1–3 month delay in the timing of maximum LAI. This bias was reduced in CLM3-CN where it was most significant.
- Both models overestimate woody biomass in the Amazon Basin. Carbon comparisons with Malhi *et al.* (in press) suggest too much allocation to wood. Allocation in CLM3-CN was adjusted to reduce this bias.
- The models differed by a factor of two in annual carbon sinks. Both results are compatible with atmospheric budgets given other uncertainties.

# Biases and Weaknesses Exposed by the C-LAMP Analysis

- Both models underestimated the amplitude of the seasonal cycle of  $\text{CO}_2$  in the northern hemisphere. Adjustment of the  $Q_{10}$  for heterotrophic respiration from 2.0 to 1.5 in CLM3-CASA' reduces this bias. Adoption of the same  $Q_{10}$  formulation, in place of Lloyd & Taylor, reduces this bias in CLM3-CN. The  $Q_{10}$  for maintenance respiration in CLM3-CN was also reduced from 2.0 to 1.5.



# Biases and Weaknesses Exposed by the C-LAMP Analysis

- Both models underestimated the amplitude of the seasonal cycle of CO<sub>2</sub> in the northern hemisphere. Adjustment of the  $Q_{10}$  for heterotrophic respiration from 2.0 to 1.5 in CLM3-CASA' reduces this bias. Adoption of the same  $Q_{10}$  formulation, in place of Lloyd & Taylor, reduces this bias in CLM3-CN. The  $Q_{10}$  for maintenance respiration in CLM3-CN was also reduced from 2.0 to 1.5.
- CLM3-CN seasonal cycle was out of phase with observations. A new day-length control on photosynthesis mechanism mitigates this bias in CLM3-CN.

C-LAMP Paper in Press in *Global Change Biology*

Global Change Biology (2009), doi: 10.1111/j.1365-2486.2009.01912.x

## Systematic assessment of terrestrial biogeochemistry in coupled climate–carbon models

JAMES T. RANDERSON\*, FORREST M. HOFFMAN†, PETER E. THORNTON‡§, NATALIE M. MAHOWALD¶, KEITH LINDSAY‡, YEN-HUEI LEE‡, CYNTHIA D. NEVISON\*||, SCOTT C. DONEY\*\*, GORDON BONAN‡, RETO STÖCKLI††‡‡, CURTIS COVEY§§, STEVEN W. RUNNING¶¶ and INEZ Y. FUNG|||

\*Department of Earth System Science, Croul Hall, University of California, Irvine, CA 92697, USA, †Oak Ridge National Laboratory, Computational Earth Sciences Group, PO Box 2008, Oak Ridge, TN 37831, USA, ‡Climate and Global Dynamics, National Center for Atmospheric Research, PO Box 3000, Boulder, CO 80307, USA, §Oak Ridge National Laboratory, Environmental Sciences Division, PO Box 2008, Oak Ridge, TN 37831, USA, ¶Department of Earth and Atmospheric Sciences, 2140 Snee Hall, Cornell University, Ithaca, NY 14850, USA, ||Institute for Arctic and Alpine Research (INSTAAR), University of Colorado, Boulder, CO 80309, USA, \*\*Department of Marine Chemistry and Geochemistry, MS 25, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA, ††Department of Atmospheric Sciences, Colorado State University, Ft Collins, CO 80523, USA, ‡‡MeteoSwiss, Climate Service, Federal Office of Meteorology and Climatology, CH-8044 Zurich, Switzerland, §§Program for Climate Model Diagnosis and Intercomparison, 7000 East Avenue, Bldg. 170, L-103, Livermore, CA 94550-9234, USA, ¶¶Numerical Terradynamic Simulation Group, College of Forestry & Conservation, University of Montana, Missoula, MT 59812, USA, |||Department of Earth and Planetary Science and Department of Environmental Science, Policy, and Management, 307 McCone, Mail Code 4767, University of California, Berkeley, CA 94720, USA

## Abstract

With representation of the global carbon cycle becoming increasingly complex in climate models, it is important to develop ways to quantitatively evaluate model performance against *in situ* and remote sensing observations. Here we present a systematic framework, the Carbon-Land Model Intercomparison Project (C-LAMP), for assessing terrestrial biogeochemistry models coupled to climate models using observations that span a wide range of temporal and spatial scales. As an example of the value of such



## Recent Progress

- C-LAMP drove the development of model improvements in the terrestrial biogeochemistry models for the [Community Land Model version 4 \(CLM4\)](#).
- Subsequent C-LAMP analyses of six model configurations using CLM3.6 (a pre-release version of CLM4) with CASA' and CN demonstrated better performance by CN.
- Therefore, the CLM4 release will include CN. That configuration will probably be called CLM4-BGC.
- CLM4-BGC will be part of the [Community Climate System Model version 4 \(CCSM4\)](#), which may be called the Community Earth System Model (CESM). This model will be used for IPCC AR5 simulations.
- The physical models for CCSM4 are expected to be finalized before the end of 2009, and the full ESM configuration will follow within six months.


# iLAMB – International Land Model Benchmarking

- We believe that C-LAMP should serve as a prototype for a wider international benchmarking activity, as we have discussed at this meeting.
- Needed are
  - ① a well-crafted protocol that exercises model capabilities for simulating energy, water, and biogeochemical cycles;
  - ② model output data and metadata standards to simplify subsequent analyses;
  - ③ best-available forcing data sets; and
  - ④ best-available observational data sets and diagnostics.
- Follow-on discussions should be held at iLEAPS in Melbourne in August and ICDC8 in Jena in September.
- We could finalize the protocol, output standards, and relationship to AR5 simulations at a meeting in Spring 2010 at Irvine, California or Oak Ridge, Tennessee.

Thank you!

Questions?

More Discussion?



Contact: Forrest Hoffman ([forrest@climatemodeling.org](mailto:forrest@climatemodeling.org))