


The Carbon-Land Model Intercomparison Project (C-LAMP): A Prototype for Coupled Biosphere-Atmosphere Model Benchmarking for the IPCC Fifth Assessment Report (AR5)

Forrest M. Hoffman¹, James T. Randerson², Inez Y. Fung³
Peter E. Thornton¹, Natalie M. Mahowald⁴,
Gordon B. Bonan⁵, and Steven W. Running⁶

¹Oak Ridge National Laboratory, ²University of California-Irvine, ³University of California-Berkeley, ⁴Cornell University, ⁵University of Colorado-Boulder, and ⁶University of Montana

6th GEWEX/2nd iLEAPS Parallel Science Conferences
27 August 2009 • Melbourne, Australia

- The **Carbon-Land Model Intercomparison Project (C-LAMP)** began as a **CCSM Biogeochemistry Working Group** project to assess model capabilities in the coupled climate system and to explore processes important for inclusion in the CCSM4 Earth System Model for use in the IPCC Fifth Assessment Report (AR5).
- Unlike traditional MIPs, C-LAMP was designed to confront models with best-available observational datasets, develop metrics for evaluation of biosphere models, and build a general-purpose biogeochemistry diagnostics package for model evaluation.



- C-LAMP is a Biogeochemistry Subproject of the **Computational Climate Science End Station** (Warren Washington, PI), a U.S. Dept. of Energy INCITE Project.
- Models were initially run on the Cray X1E vector supercomputer in ORNL's **National Center for Computational Sciences (NCCS)**.
Cray X1E (phoenix)



1024 processors (MSPs), 2048 GB memory, and 18.08 TFlop/s peak
DECOMMISSIONED September 30, 2008

Present Jaguar: 250 TFlop/s



New Jaguar: Second Fastest in the World at 1.059 PFlop/s



JAGUAR

**World's Most Powerful Computer.
For Science!**

"The Jaguar system at ORNL provides immense computing power in a balanced, stable system that is allowing scientists and engineers to tackle some of the world's most challenging problems."
—2008, Kelvin Droegemeier, Meteorology Professor, University of Oklahoma.

Model Configurations

- Biosphere models coupled to the Community Climate System Model version 3.1
 - **CLM3-CASA'** — Carnegie/Ames/Stanford Approach Model previously run in CSM1.4 (Fung)
 - **CLM3-CN** — coupled carbon and nitrogen cycles based on the Biome-BGC model (Thornton)
 - **LSX-IBIS** — Integrated Biosphere Simulator from U. Wisconsin previously run in PCTM (Thompson)
- Because LSX-IBIS is not coupled to the CLM3 biophysics and was not a candidate for inclusion in CCSM4, only CLM3-CASA' and CLM3-CN were evaluated in C-LAMP.
- CCSM3.1 partially coupled ("I" & "F" configurations) run at T42 resolution ($\sim 2.8^\circ \times 2.8^\circ$), spectral Eulerian dycore, $1^\circ \times 0.27^\circ$ – 0.53° ocean & sea ice data models (T42gx1v3).

C-LAMP Protocol Overview

- **Experiment 1:** Models forced with an improved NCEP/NCAR reanalysis climate data set (Qian, *et al.* 2006) to examine the influence of climate variability, prescribed atmospheric CO₂, and land cover change on terrestrial carbon fluxes during the 20th century (specifically 1948–2004).
- **Experiment 2:** Models coupled with an active atmosphere (CAM3), prescribed atmospheric CO₂, prescribed sea surface temperatures and ocean carbon fluxes to examine the effect of a coupled biosphere-atmosphere for carbon fluxes and climate during the 20th century.
- All the forcing and observational datasets are being shared, and model results are available through the Earth System Grid (ESG), just like for CMIP3 (the IPCC AR4 model results).
- Experimental protocol, output fields, and metrics are available at <http://www.climatemodeling.org/c-lamp/>

Offline Forcing with NCEP/NCAR Reanalysis

Exp.	Description	Time Period
1.1	Spin Up	~4,000 y
1.2	Control	1798–2004
1.3	Varying climate	1948–2004
1.4	Varying climate, CO ₂ , and N deposition	1798–2004
1.5	Varying climate, CO ₂ , N deposition and land use	1798–2004
1.6	Free Air CO ₂ Enrichment (FACE) Control	1997–2100
1.7	Free Air CO ₂ Enrichment (FACE) Transient	1997–2100

Coupled Land-Atmosphere Forcing with Hadley SSTs

Exp.	Description	Time Period
2.1	Spin Up	~2,600 y
2.2	Control	1800–2004
2.3	Varying climate	1800–2004
2.4	Varying climate, CO ₂ , and N deposition	1800–2004
2.5	Varying climate, CO ₂ , N deposition and land use	1800–2004
2.6	Varying climate, CO ₂ , N deposition, seasonal FFE	1800–2004

All but the land use experiments were run with CCSM3.1
using CLM3-CASA' and CLM3-CN biogeochemistry models
yielding >16,000 y and ~50 TB

C-LAMP Common Model Output - Mozilla Firefox

file Edit View History Bookmarks Tools Help

http://www.climate modeling.org/c-lamp/protocol/model_output.php

C-LAMP Common Model Output

While all models participating in the Carbon Land Model intercomparison Project (C-LAMP) will output their own "native" fields, a common set of fields is needed to facilitate head-to-head comparison of the models to each other and to available observational datasets. Model results transmitted to the [Earth System Grid](#) for redistribution to the community will use common field names, netCDF long names, [CF Standard Names](#) and units. Contained below is a table of the common output fields required for the C-LAMP and consistent with the metadata conventions used for [CMIP3](#), formerly called the IPCC 4th Assessment Model Output database. Corrections and suggestions are solicited on this information. Software is available for rewriting model output into netCDF files following the [Climate and Forecast \(CF\) Metadata Convention](#).

Version 2.1 - Aug 30, 2008

Atmospheric forcing				
Variable Name	Long Name and CF Standard Name	Units	Comment	Statistics
husf	Specific humidity at atmospheric forcing height specific_humidity†	kg kg-1		MHM, MHS, MM
prra	Rainfall precipitation flux rainfall_flux†	kg m-2 s-1	Rainfall includes all liquid types (rain, large-scale, convective, etc.)	MHM, MHS, MM
prsn†	Snowfall precipitation flux snowfall_flux†	kg m-2 s-1	Snowfall includes all frozen types (snow, hail, ice, etc.)	MHM, MHS, MM

Biogeochemistry				
Variable Name	Long Name and CF Standard Name	Units	Comment	Statistics
agbc*	Above-ground biomass carbon above_ground_biomass_carbon_content	kg m-2	Total carbon content in above-ground live and dead carbon pool(s)	MM
aglbc*	Above-ground live biomass carbon above_ground_live_biomass_carbon_content	kg m-2	Total carbon content in above-ground live carbon pool(s)	MM
agnpp	Above-ground net primary production above_ground_net_primary_productivity_of_carbon	kg m-2 s-1	Component of net primary production attributable to above-ground live biomass	MM
ar	Autotrophic respiration autotrophic_respiration_of_carbon alias(es): plant_respiration_carbon_flux	kg m-2 s-1	Sum of maintenance respiration and growth respiration of vegetation	MHM, MHS, MM
bco	Biogenic carbon monoxide flux biogenic_carbon_monoxide_flux	kg m-2 s-1	Total biogenic carbon monoxide flux out of biosphere	MM

Done

C-LAMP Performance Metrics and Diagnostics

- An evolving document on metrics for model evaluation is available at <http://www.climatemodeling.org/c-lamp/>
- Each model is scored with respect to its performance on various output fields compared with best-available observational datasets.
- Examples include:
 - leaf area index (LAI): comparison of phase and spatial distribution using MODIS
 - net primary production (NPP): comparison with EMDI and correlation with MODIS
 - CO₂ seasonal cycle: comparison with NOAA/Globalview flask sites after combining fluxes with impulse response functions from TRANSCOM
 - regional carbon stocks (Saatchi *et al.*, 2006; Batjes, 2006)
 - carbon and energy fluxes (Fluxnet sites)
 - other transient dynamics: β factor, fire emissions

Score Sheet for CLAMP - Mozilla Firefox

File Edit View History Bookmarks Tools Help

http://www.climate modeling.org/c-lamp/results/diagnostics/CN_vs_CN/ Google

C-LAMP Score Sheet for Biogeochemical Model Evaluation


Metric	Metric components	Observations & comparison protocol	Model CASA'	Model CN	Score (points)		
					Possible	CASA'	CN
LAI	MODIS Phase	global map	global map model vs obs	global map model vs obs	6.00	5.11	4.24
	MODIS Maximum	global map	global map model vs obs	global map model vs obs	5.00	4.60	4.26
	MODIS Mean	land class obs land class model global map	model vs obs table global map model vs obs	model vs obs table global map model vs obs	4.00	3.75	3.53
NPP	EMDI NPP observations	Class A table	table scatter plot	table scatter plot	1.00	0.68	0.73
		Class B table	table scatter plot	table scatter plot	1.00	0.83	0.82
	EMDI NPP normalized by PPT	Class A histogram	Class A histogram	Class A histogram	2.00	1.50	1.74
		Class B histogram	Class B histogram	Class B histogram	2.00	1.51	1.65
	Correlation with MODIS	global map	model map model vs obs	model map model vs obs	2.00	1.64	1.44
Correlation with MODIS-zonal mean	zonal mean obs	zonal mean model vs obs plot	zonal mean model vs obs plot	2.00	1.88	1.84	
CO ₂ Seasonal Cycle — Comparison with Globalview phase and amplitude	60°N–90°N	—	—	—	6.00	4.11	2.77
	30°N–60°N	—	—	—	6.00	4.23	3.23
	0°N–30°N	—	—	—	3.00	2.07	1.71
Energy and C Fluxes from Fluxnet	NEE	—	—	—	—	—	—
	Net radiation	—	—	—	—	—	—
	Latent heat	line plot	model vs obs	model vs obs	—	—	—
Energy and C Fluxes from Ameriflux	Sensible heat	—	—	—	—	—	—
	NEE	—	—	—	6.00	2.46	2.13
	Shortwave Incoming	—	—	—	—	—	—
	Latent heat	line plot	model vs obs timeseries plot	model vs obs timeseries plot	9.00	6.38	6.39
	Sensible heat	—	—	—	9.00	4.90	4.64
	GPP	—	—	—	6.00	3.39	3.46
	ER	—	—	—	—	—	—
Aboveground live biomass in South	obs amazon	model amazon	amazon map	10.00	5.20	4.00	

Done

Score Sheet for CLAMP - Mozilla Firefox								
File Edit View History Bookmarks Tools Help								
http://www.climate modeling.org/c-lamp/results/diagnostics/CN_vs_C/ Google								
CO ₂ Seasonal Cycle — Comparison with Globalview phase and amplitude	60°N–90°N	—	—	—	6.00	4.11	2.77	
	30°N–60°N	—	—	—	6.00	4.23	3.23	
	0°N–30°N	—	—	—	3.00	2.07	1.71	
Energy and C Fluxes from Fluxnet	NEE	—	—	—	—	—	—	
	Net radiation	—	—	—	—	—	—	
	Latent heat	line plot	model vs obs	model vs obs	—	—	—	
	Sensible heat	—	—	—	—	—	—	
Energy and C Fluxes from Ameriflux	NEE	—	—	—	6.00	2.46	2.13	
	Shortwave Incoming	—	—	—	—	—	—	
	Latent heat	line plot	model vs obs timeseries plot	model vs obs timeseries plot	9.00	6.38	6.39	
	Sensible heat	—	—	—	9.00	4.90	4.64	
	GPP	—	—	—	6.00	3.39	3.46	
	ER	—	—	—	—	—	—	
Carbon Stocks and Transient Dynamics	Aboveground live biomass in South America	obs amazon	model amazon model vs obs	amazon map model vs obs	10.00	5.28	4.99	
	Aboveground live biomass within Amazon Basin (sum within Legal Amazon)	mask obs masked 68.90 (Pg C)	model masked model vs obs 198.87 (Pg C)	model masked model vs obs 160.61 (Pg C)	—	—	—	
	NPP Stimulation from elevated CO ₂	—	FACE Site table biome table	FACE Site table biome table	10.00	7.87	4.11	
	Interannual variability of global carbon fluxes - comparison with TRANSCOM	—	—	—	5.00	3.55	3.00	
	Turnover times and pool sizes	—	Leaf Wood Fine Root Litter Coarse Woody Debris Soil	Leaf Wood Fine Root Litter Coarse Woody Debris Soil	—	—	—	
	Carbon Sinks (1990–2004)	—	biome mean biome total	biome mean biome total	—	—	—	
	Fire Variability (1997–2004)	—	—	global spatial comparison temporal dynamics	5.00	—	1.70	
	Total Score				100.00	65.74	58.38	


Done

- Comparisons with field observations include net primary production (NPP) from the Ecosystem Model-Data Intercomparison (EMDI).
- Measurements were performed in different ways, at different times, and by different groups for a limited number of field sites.
- Shown here are comparisons of NPP with EMDI Class A observations (Figures a and b) and Class B observations (Figures c and d).




Data provided by NASA Distributed Active Archive Center (DAAC) at ORNL


- Comparisons with satellite “modeled observations” must be made carefully because of high uncertainty.
- This comparison with MODIS leaf area index (LAI) focuses on the month of maximum LAI (phase), a measurement with less uncertainty than the “observed” LAI values.
- C-LAMP accounts for this uncertainty by weighting scores accordingly.
- CLM-CASA' scored 5.1/6.0 while CLM-CN scored 4.2/6.0 for this metric.




- MODIS net primary production (NPP) “observations” have higher uncertainty.
- Comparison with MODIS NPP focuses on correlation of spatial patterns.
- CLM-CASA' scored 1.6/2.0 while CLM-CN scored 1.4/2.0.




- Comparisons with Globalview flask sites are made by combining model fluxes with impulse response functions from TRANSCOM.
- Shown are the annual cycles of atmospheric CO₂ at (a) Mould Bay, Canada (76°N), (b) Storhofdi, Iceland (63°N), (c) Carr, Colorado (41°N), (d) Azores Islands (39°N), (e) Sand Island, Midway (28°N), and (f) Kumakahi, Hawaii (20°N).
- CLM-CASA' scored 10.4/15.0 while CLM-CN scored 7.7/15.0 for this metric.



- Estimates of carbon stocks are very difficult to obtain.
- This comparison with estimates of aboveground live biomass in the Amazon by Saatchi *et al.* (2006) shows that both models are too high by about a factor of 2.
- Using a score based on normalized cell-by-cell differences, CLM-CASA' scored 5.3/10.0 while CLM-CN scored 5.0/10.0.






- Comparisons with AmeriFlux eddy correlation CO₂ flux tower sites include net ecosystem exchange (NEE), gross primary production (GPP), respiration, shortwave incoming radiation, and latent and sensible heat.
- Shown here is a comparison of model estimates with eddy covariance measurements from Sylvania Wilderness, Harvard Forest, and Walker Branch.
- Used are the consistent Level 4 data produced by Dario P. and Markus R.



Data provided by ORNL Carbon Dioxide Information Analysis Center (CDIAC).

- Additional field measurement comparisons include the Free Air CO₂ Enrichment (FACE) results, including the ORNL site.
- The Norby *et al.* (2005) synthesis of four FACE site observations suggested “response of forest NPP to elevated [CO₂] is highly conserved across a broad range of productivity, with a stimulation at the median of $23 \pm 2\%$.”
- A C-LAMP experiment was added to test this result by increasing [CO₂] to 550 ppmv in 1997.





Site Name	Lon (°E)	Lat (°N)	Observations		CASA'			CN		
			NPP↑	β_L	NPP↑	β_L	Score	NPP↑	β_L	Score
Duke	-79.08	35.97	28.0%	0.69	16.4%	0.41	0.26	6.2%	0.15	0.65
Aspen	-89.62	45.67	35.2%	0.87	15.6%	0.39	0.39	12.4%	0.31	0.48
ORNL	-84.33	35.90	23.9%	0.59	17.3%	0.43	0.16	5.2%	0.13	0.64
POP-Euro	11.80	42.37	21.8%	0.54	20.0%	0.49	0.04	5.7%	0.14	0.59
4 site mean			27.2%	0.67	17.3%	0.43		7.4%	0.18	
Total M Score							0.79	0.41		

But! Norby is now reporting reduced NPP enhancement at the ORNL FACE site due probably to N limitation!

C-LAMP Score Sheet for CLM3-CASA' and CLM3-CN

Metric	Metric components	Uncertainty of obs.	Scaling mismatch	Total score	Sub-score	CASA'	CN
LAI	Matching MODIS observations			15.0		13.5	12.0
	• Phase (assessed using the month of maximum LAI)	Low	Low		6.0	5.1	4.2
	• Maximum (derived separately for major biome classes)	Moderate	Low		5.0	4.6	4.3
	• Mean (derived separately for major biome classes)	Moderate	Low		4.0	3.8	3.5
NPP	Comparisons with field observations and satellite products			10.0		8.0	8.2
	• Matching EMDI Net Primary Production observations	High	High		2.0	1.5	1.6
	• EMDI comparison, normalized by precipitation	Moderate	Moderate		4.0	3.0	3.4
	• Correlation with MODIS (r^2)	High	Low		2.0	1.6	1.4
	• Latitudinal profile comparison with MODIS (r^2)	High	Low		2.0	1.9	1.8
CO ₂ annual cycle	Matching phase and amplitude at Globalview flash sites			15.0		10.4	7.7
	• 60°–90°N	Low	Low		6.0	4.1	2.8
	• 30°–60°N	Low	Low		6.0	4.2	3.2
	• 0°–30°N	Moderate	Low		3.0	2.1	1.7
Energy & CO ₂ fluxes	Matching eddy covariance monthly mean observations			30.0		17.2	16.6
	• Net ecosystem exchange	Low	High		6.0	2.5	2.1
	• Gross primary production	Moderate	Moderate		6.0	3.4	3.5
	• Latent heat	Low	Moderate		9.0	6.4	6.4
	• Sensible heat	Low	Moderate		9.0	4.9	4.6
Transient dynamics	Evaluating model processes that regulate carbon exchange on decadal to century timescales			30.0		16.8	13.8
	• Aboveground live biomass within the Amazon Basin	Moderate	Moderate		10.0	5.3	5.0
	• Sensitivity of NPP to elevated levels of CO ₂ : comparison to temperate forest FACE sites	Low	Moderate		10.0	7.9	4.1
	• Interannual variability of global carbon fluxes: comparison with TRANSCOM	High	Low		5.0	3.6	3.0
	• Regional and global fire emissions: comparison to GFEDv2	High	Low		5.0	0.0	1.7
	Total:				100.0		65.9

Earth System Grid (ESG) Node at ORNL for C-LAMP

C-LAMP Model Data - Mozilla Firefox

File Edit View Go Bookmarks Tools Help

https://esg2.ornl.gov:8443/

C-LAMP Model Data

Home Data About ESG Login

Contact ESG

Scientific Discovery through Advanced Computing

Welcome

Welcome to the CCES C-LAMP data portal. If you are new to this site, please review the help pages:

- Registration
- Searching
- Browsing and Downloading Data
- Downloading from FTP

Data Search

Search Dataset metadata for:

Example: nri, cccma

[Advanced Search](#)

Browse Dataset Catalogs

- CCSM Carbon LAnd Model Intercomparison Project (C-LAMP)

CCES C-LAMP Portal Collaborators

PCMDI

Home | Data | About ESG | Login

Login Status: Not logged in.

© 2004, UCAR. All rights reserved.
Portions © 2004, The Regents of the University of California. All rights reserved.
[Privacy & Security Notices](#)

Done esg2.ornl.gov:8443

Biases and Weaknesses Exposed by the C-LAMP Analysis

- Both models had a low LAI bias in boreal and arctic regions. This bias was partially eliminated by a new hydrology model capturing freeze-thaw dynamics.
- Both models had a 1–3 month delay in the timing of maximum LAI. This bias was reduced in CLM3-CN where it was most significant.
- Both models overestimate woody biomass in the Amazon Basin. Carbon comparisons with Malhi *et al.* (in press) suggest too much allocation to wood. Allocation in CLM3-CN was adjusted to reduce this bias.
- The models differed by a factor of two in annual carbon sinks. Both results are compatible with atmospheric budgets given other uncertainties.

Biases and Weaknesses Exposed by the C-LAMP Analysis

- Both models underestimated the amplitude of the seasonal cycle of CO_2 in the northern hemisphere. Adjustment of the Q_{10} for heterotrophic respiration from 2.0 to 1.5 in CLM3-CASA' reduces this bias. Adoption of the same Q_{10} formulation, in place of Lloyd & Taylor, reduces this bias in CLM3-CN. The Q_{10} for maintenance respiration in CLM3-CN was also reduced from 2.0 to 1.5.
- CLM3-CN seasonal cycle was out of phase with observations. A new day-length control on photosynthesis mechanism mitigates this bias in CLM3-CN.

C-LAMP Paper in *Global Change Biology*

Global Change Biology (2009), doi: 10.1111/j.1365-2486.2009.01912.x

Systematic assessment of terrestrial biogeochemistry in coupled climate–carbon models

JAMES T. RANDERSON*, FORREST M. HOFFMAN†, PETER E. THORNTON‡§, NATALIE M. MAHOWALD¶, KEITH LINDSAY‡, YEN-HUEI LEE‡, CYNTHIA D. NEVISON*||, SCOTT C. DONEY**, GORDON BONAN‡, RETO STÖCKLI††‡‡, CURTIS COVEY§§, STEVEN W. RUNNING¶¶ and INEZ Y. FUNG|||

*Department of Earth System Science, Croul Hall, University of California, Irvine, CA 92697, USA, †Oak Ridge National Laboratory, Computational Earth Sciences Group, PO Box 2008, Oak Ridge, TN 37831, USA, ‡Climate and Global Dynamics, National Center for Atmospheric Research, PO Box 3000, Boulder, CO 80307, USA, §Oak Ridge National Laboratory, Environmental Sciences Division, PO Box 2008, Oak Ridge, TN 37831, USA, ¶Department of Earth and Atmospheric Sciences, 2140 Snee Hall, Cornell University, Ithaca, NY 14850, USA, ||Institute for Arctic and Alpine Research (INSTAAR), University of Colorado, Boulder, CO 80309, USA, **Department of Marine Chemistry and Geochemistry, MS 25, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA, ††Department of Atmospheric Sciences, Colorado State University, Ft Collins, CO 80523, USA, ‡‡MeteoSwiss, Climate Service, Federal Office of Meteorology and Climatology, CH-8044 Zurich, Switzerland, §§Program for Climate Model Diagnosis and Intercomparison, 7000 East Avenue, Bldg. 170, L-103, Livermore, CA 94550-9234, USA, ¶¶Numerical Terradynamic Simulation Group, College of Forestry & Conservation, University of Montana, Missoula, MT 59812, USA, |||Department of Earth and Planetary Science and Department of Environmental Science, Policy, and Management, 307 McCone, Mail Code 4767, University of California, Berkeley, CA 94720, USA

Abstract

With representation of the global carbon cycle becoming increasingly complex in climate models, it is important to develop ways to quantitatively evaluate model performance against *in situ* and remote sensing observations. Here we present a systematic framework, the Carbon-Land Model Intercomparison Project (C-LAMP), for assessing terrestrial biogeochemistry models coupled to climate models using observations that span a wide range of temporal and spatial scales. As an example of the value of such



Recent Progress

- C-LAMP helped drive the development of model improvements in the terrestrial biogeochemistry models for the [Community Land Model version 4 \(CLM4\)](#).
- Subsequent C-LAMP analyses of six model configurations using CLM3.6 (a pre-release version of CLM4) with CASA' and CN demonstrated much improved performance by CN.
- It is now recognized that physical model changes must be tested using C-LAMP to ensure that these changes do not have negative impacts on biogeochemistry model performance.
- While our recent proposal to deploy C-LAMP as a web service was not funded, we are sharing the data and diagnostics package for others to use (e.g., Jena's JEDI model) and hoping to incorporate additional metrics over time.
- **Next: N-LAMP** — develop a strategy for benchmarking the nitrogen cycle in land surface models.

C-LAMP, C⁴MIP, and iLEAPS

- C-LAMP Experiment 2 is patterned after C⁴MIP (Coupled Climate-Carbon Cycle Model Intercomparison Project, <http://www.c4mip.cnrs-gif.fr/>) Phase 1.
- At the October 2006 C⁴MIP Workshop at the UK Met Office in Exeter, there was strong interest in Experiment 1 and validation experiments using Fluxnet observations.
- At the Marie Curie/iLEAPS Workshop in Hyères in November 2008, a number of modeling groups expressed interest in consistent model validation and model-data comparisons for their coupled biosphere models. See write up in iLEAPS Newsletter number 7.
- A QUEST/GLASS model benchmarking workshop was held in June 2009 at the University of Exeter where a strategy was discussed for combining Australian, European, and U.S. efforts toward a truly international benchmarking system.


C-LAMP + ILAMB + ...

- We believe that C-LAMP and ILAMB should serve as a prototype for a wider international benchmarking activity, the results of which could contribute to AR5.
- Needed are
 - 1 a well-crafted protocol that exercises model capabilities for simulating energy, water, and biogeochemical cycles;
 - 2 model output data and metadata standards to simplify subsequent analyses;
 - 3 best-available forcing data sets; and
 - 4 best-available observational data sets and diagnostics.
- Follow-on discussions should be held by carbon cycle researchers at ICDC8 in Jena in September.
- We could finalize the protocol, output standards, metrics and diagnostics, and relationship to AR5 simulations at a meeting in Spring 2010 in the U.S.

Thank you!

Questions?

More Discussion?



Contact: Forrest Hoffman (forrest@climatemodeling.org)